

Determination of the ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ asymptotic normalization coefficients (nuclear vertex constants) and their application for extrapolation of the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factors to the solar energy region

S.B. Igamov, Q.I. Tursunmahatov and R. Yarmukhamedov *

January 6, 2010

Institute of Nuclear Physics, Uzbekistan Academy of Sciences, 100214 Tashkent, Uzbekistan

Abstract

A new analysis of the precise experimental astrophysical S factors for the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction [B.S. Nara Singh et al., Phys.Rev.Lett. **93**, 262503 (2004); D. Bemmerer et al., Phys.Rev.Lett. **97**, 122502 (2006); F.Confortola et al., Phys.Rev.C **75**, 065803 (2007) and T.A.D.Brown et al., Phys.Rev. C **76**, 055801 (2007)] populating to the ground and first excited states of ${}^7\text{Be}$ is carried out based on the modified two-body potential approach in which the direct astrophysical S factor, $S_{34}(E)$, is expressed in terms of the asymptotic normalization constants for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$. The Woods-Saxon potential form is used for the bound $(\alpha + {}^3\text{He})$ -state and the α - ${}^3\text{He}$ -scattering wave functions. New estimates are obtained for the "indirectly determined" values of the asymptotic normalization constants (the nuclear vertex constants) for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(g.s.)$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429\text{MeV})$ as well as the astrophysical S factors $S_{34}(E)$ at $E \leq 90$ keV, including $E=0$. The values of asymptotic normalization constants have been used for getting information about the α -particle spectroscopic factors for the mirror (${}^7\text{Li}$ - ${}^7\text{Be}$)-pair.

PACS: 25.40.Lw; 26.35.+c

1 Introduction

The ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction is one of the critical links in the ${}^7\text{Be}$ and ${}^8\text{B}$ branches of the pp -chain of solar hydrogen burning [1–3]. The total capture rate determined by processes of this chain is sensitive to the cross section $\sigma_{34}(E)$ (or the astrophysical S factor $S_{34}(E)$) for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction and predicted neutrino rate varies as $[S_{34}(0)]^{0.8}$ [2, 3].

*Corresponding author, E-mail: rakhim@inp.uz

Despite the impressive improvements in our understanding of the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction made in the past decades (see Refs [4–10] and references therein), however, some ambiguities connected with both the extrapolation of the measured cross sections for the aforesaid reaction to the solar energy region and the theoretical predictions for $\sigma_{34}(E)$ (or $S_{34}(E)$) still exist and they may influence the predictions of the standard solar model [2, 3] .

Experimentally, there are two types of data for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction at extremely low energies: i) six measurements based on detecting of γ -rays capture (see [4] and references therein) from which the astrophysical S factor $S_{34}(0)$ extracted by the authors of those works changes within the range $0.47 \leq S_{34}(0) \leq 0.58$ keV b, which yield a weighted mean of $S_{34}(0) = 0.507 \pm 0.016$ keV b [4], and ii) five measurements based on detecting of ${}^7\text{Be}$ (see [4] references therein as well as [6–10]) from which $S_{34}(0)$ extracted by the authors of these works changes within the range $0.53 \leq S_{34} \leq 0.63$ keV b, which yield weighted means of $S_{34}(0) = 0.572 \pm 0.026$ keV b [4], $S_{34}(0) = 0.53$ keV b [6], $S_{34}(0) = 0.547 \pm 0.017$ keV b [7, 8], and $S_{34}(0) = 0.560 \pm 0.017$ keV b [9] and $S_{34}(0) = 0.595 \pm 0.018$ and 0.596 ± 0.021 keV b [10]. All of these measured data have a similar energy dependence for the astrophysical S factors $S_{34}(E)$ but the extrapolation of each of the measured data from the observed energy ranges to low experimentally inaccessible energy regions, including $E=0$, gives a value of $S_{34}(0)$ with an uncertainty exceeding noticeably the experimental one. The recent aforesaid values of $S_{34}(0)$ recommended in Refs.[6–9] and [10] have nevertheless been obtained from the analysis of the precisely measured data for $S_{34}^{exp}(E)$ by means of the artificial renormalization of the energy dependence of the R-matrix calculation [11] and of the resonating-group method calculation [12] for $S_{34}(E)$ to the corresponding experimental data, respectively.

The theoretical calculations of $S_{34}(0)$ performed within different methods also show considerable spread [12–17]. The aforesaid resonating-group method calculations of $S_{34}(0)$ performed in Ref.[12] show considerable sensitivity to the form of the effective NN interaction used and the estimates have been obtained within the range of $0.312 \leq S_{34}(0) \leq 0.841$ keV b. Calculations performed in microscopic single-channel ($(\alpha + {}^3\text{He})$) and two-channel ($({}^3\text{He} + \alpha)$ and $(p + {}^6\text{Li})$) cluster models gave the values of $S_{34}(0) = 0.56$ keV b [13] ($S_{34}(0) = 0.52$ keV b [14]) and of $S_{34}(0) = 0.83$ keV b [14], respectively, that is, the estimate of the value of the $S_{34}(0)$ strongly changes when the model space is expanded. Also, one would like to note calculations performed in the variational Monte-Carlo technique (VMCT) with seven-particle wave functions derived from realistic NN interaction gave $S_{34}(0) \approx 0.40$ keV b [15]. But, as it was emphasized in paper [15], serious problems occur with the normalization for the calculated astrophysical S factor $S(E)$ in respect to the experimental data. The calculations performed by the authors of Refs.[16] and [17] in the two-body potential model with different forms of the two-body potential gave the values of $S_{34}(0) = 0.516$ and about of 0.5 keV b, respectively, although different values of 1.174 in [16] and 1.0 in [17] have been used for the spectroscopic factor for the $(\alpha + {}^3\text{He})$ -configuration in ${}^7\text{Be}$. But, it should be noted that in [16, 17] a use of the normalization of the astrophysical S factor $S_{34}(E)$ in terms of the spectroscopic factors is hardly justified since the spectroscopic factor determines the probability of finding the $(\alpha + {}^3\text{He})$ -configuration in ${}^7\text{Be}$, which has been created in the reaction under consideration, and this probability is mainly determined by the radial part of the overlap function of the bound state wave functions for the α , ${}^3\text{He}$ and ${}^7\text{Be}$ nuclei in the nuclear interior, while the dominant contribution to the aforesaid

peripheral reaction comes from the surface and external regions of the nucleus ${}^7\text{Be}$ [18, 19](see, below also).

The estimation of $S_{34}(0)=0.52\pm 0.03$ keV b [19] also should be noted. The latter has been obtained within the framework of the standard two-body potential model in the assumption that the contribution from the nuclear interior ($r < r_{cut}$, $r_{cut}=4$ fm) to the amplitude is negligibly small [20, 21]. Therefore, in [19] the radial wave function for the bound (${}^3\text{He} + \alpha$) state of the nucleus ${}^7\text{Be}$ in the integrand of the radial integral (see Eq. (10) of Ref.[22] for example) was replaced by the appropriated Whittaker function and, consequently, the astrophysical S factor was directly expressed in terms of the nuclear vertex constants (NVC) for the virtual decays ${}^7\text{Be} \rightarrow \alpha + {}^3\text{He}$ (or respective the asymptotic normalization coefficients (ANC) for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$) [23, 24]. As a result, in Ref. [19], from the analysis of the experimental astrophysical S factors for the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ reactions in the energy range $180 \lesssim E \lesssim 500$ keV [25], the values of the nuclear vertex constants (NVC) for the virtual decays ${}^7\text{Be}(\text{g.s.}) \rightarrow \alpha + {}^3\text{He}$ and ${}^7\text{Be}(0.429 \text{ MeV}) \rightarrow \alpha + {}^3\text{He}$ were obtained, which were then used for calculations of the astrophysical S factors for the same reactions at $E < 180$ keV, including $E=0$. However, the experimental astrophysical S factors for the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reactions [25] used in [19] for the analysis have considerable spread. Consequently, the values of the ANC's (or NVC's) ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ and the $S_{34}(0)$ obtained in [19] may not be enough accurately associated both with the aforesaid assumption in respect to the contribution from the nuclear interior ($r < r_{cut}$) and with a presence of the spread in the experimental data used for the analysis. Therefore, determination of precise experimental values of the ANC's for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ is highly desirable since it has direct effects in the correct extrapolation of the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factor at solar energies.

Recently, a modified two-body potential approach (MTBPA) was proposed in [22] for the peripheral direct capture $A(a, \gamma)B$ reaction, which is based on the idea proposed in paper [18] that low-energy direct radiative captures of particle a by light nuclei A proceed mainly in regions well outside the range of the internuclear aA interactions. One notes that in MTBPA the direct astrophysical S factor is expressed in terms of ANC for $A + a \rightarrow B$ rather than through the spectroscopic factor for the nucleus B in the $(A + a)$ configuration as it is made within the standard two-body potential method [26, 27]. In Refs.[22, 28, 29], MTBPA has been successfully applied to the radiative proton and α -particle capture by some light nuclei. Therefore, it is of great interest to apply the MTBPA for analysis of the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction.

In this work new analysis of the highly precise experimental astrophysical S factors for the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction at extremely low energies ($\gtrsim 90$ keV) [6-10] is performed within the MTBPA [22] to obtain "indirectly determined" values both of the ANC's (the NVC's) for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$, and of $S_{34}(E)$ at $E \leq 90$ keV, including $E=0$. In this work we quantitatively show that the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction within the aforesaid energy region is mainly peripheral and one can extract ANC's for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ directly from the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction where the ambiguities inherent for the standard two-body potential model calculation of the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction, which is connected with the choice of the geometric parameters (the radius R and the diffuseness a) for the Woods-Saxon potential and the spectroscopic factors [16, 17], can be reduced in the physically acceptable limit, being within the experimental errors for the $S_{34}(E)$.

The contents of this paper are as follows. In Section 2 basic formulae of the modified two-body potential approach to the direct radiative capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction are given. There the analysis of the precise measured astrophysical S factors for the direct radiative capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction is performed (Subsections 2.2–2.4). The conclusion is given in Section 3.

2 Analysis of ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction

2.1 Basic formulae

Here we give the formulae specialized for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factor. Let us write l_f (j_f) for the relative orbital (total) angular momentum of ${}^3\text{He}$ and α -particle in nucleus ${}^7\text{Be}$ ($\alpha + {}^3\text{He}$), l_i (j_i) for the orbital (total) angular momentum of the relative motion of the colliding particles in the initial state, λ for multipole order of the electromagnetic transition, η_{Be} for the Coulomb parameter for the ${}^7\text{Be}$ ($\alpha + {}^3\text{He}$) bound state and μ for the reduced mass of the $(\alpha{}^3\text{He})$ -pair. For the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction populating to the ground and first excited ($E^*=0.429$ MeV; $J^\pi=1/2^-$) states of ${}^7\text{Be}$, the values of j_f are taken to be equal to $3/2$ and $1/2$, respectively, the value of l_f is taken to be equal to 1 as well as $l_i=0, 2$ for the $E1$ -transition and $l_i=1$ for the $E2$ -transition.

According to [22], for fixed l_f and j_f we can write the astrophysical S factor, $S_{l_f j_f}(E)$, for the peripheral direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction in the following form

$$S_{l_f j_f}(E) = C_{l_f j_f}^2 \mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)}). \quad (1)$$

Then, the total astrophysical S factor for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.}+0.429 \text{ MeV})$ reaction is given by

$$S_{34}(E) = \sum_{l_f=1; j_f=1/2, 3/2} S_{l_f j_f}(E) = C_{1\ 3/2}^2 \mathcal{R}_{1\ 3/2}(E, C_{1\ 3/2}^{(sp)}) + C_{1\ 1/2}^2 \mathcal{R}_{1\ 1/2}(E, C_{1\ 1/2}^{(sp)}) \quad (2)$$

$$= C_{1\ 3/2}^2 [\mathcal{R}_{1\ 3/2}(E, C_{1\ 3/2}^{(sp)}) + \lambda_C \mathcal{R}_{1\ 1/2}(E, C_{1\ 1/2}^{(sp)})] \quad (3)$$

$$= C_{1\ 3/2}^2 \mathcal{R}_{1\ 3/2}(E, C_{1\ 3/2}^{(sp)}) [1 + R(E)] \quad (4)$$

in which $R(E) = S_{1\ 1/2}(E)/S_{1\ 3/2}(E) = C_{1\ 1/2}^2 \mathcal{R}_{1\ 1/2}(E, C_{1\ 1/2}^{(sp)}) / C_{1\ 3/2}^2 \mathcal{R}_{1\ 3/2}(E, C_{1\ 3/2}^{(sp)})$ is a branching ratio and $\lambda_C = (C_{1\ 1/2}/C_{1\ 3/2})^2$.

Here, $C_{l_f j_f}$ is the ANC for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$, which determines the amplitude of the tail of the ${}^7\text{Be}$ nucleus bound state wave function in the $(\alpha + {}^3\text{He})$ -channel and is related to the spectroscopic factor $Z_{l_f j_f}$ for the $(\alpha + {}^3\text{He})$ -configuration with the quantum numbers l_f and j_f in the ${}^7\text{Be}$ nucleus by the equation $C_{l_f j_f} = Z_{l_f j_f}^{1/2} C_{l_f j_f}^{(sp)}$ [23], and $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)}) = \tilde{S}_{l_f j_f}(E) / (C_{l_f j_f}^{(sp)})^2$, where $\tilde{S}_{l_f j_f}(E) = \sum_{\lambda} \tilde{S}_{l_f j_f \lambda}(E)$ is the single-particle astrophysical S factor [5] and $C_{l_f j_f}^{(sp)}$ is the single-particle ANC, which determines the amplitude of the tail of the single-particle wave function of the bound $(\alpha + {}^3\text{He})$ state ($\varphi_{l_f j_f}(r)$). One notes that the dependence of the function $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ on the free parameter $C_{l_f j_f}^{(sp)}$ enters through the single-particle wave function

$\varphi_{l_f j_f}(r; C_{l_f j_f}^{(sp)}) (\equiv \varphi_{l_f j_f}(r))$, and the single-particle ANC $C_{l_f j_f}^{(sp)}$ in turn is itself a function of the geometric parameters of R and a ($C_{l_f j_f}^{(sp)} = C_{l_f j_f}^{(sp)}(R, a)$) [30] (see Ref.[22] also). One notes that the function $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ contains the radial integral given by Eq. (10) of Ref.[22] in which a contribution from the region of the nuclear interior ($0 \leq r \leq r_{cut}$) is also taken into account.

In Eqs. (1)–(4) the ANC's are unique unknown. As far their available values obtained by authors of Refs. [11, 15, 19], they differ noticeably from each other (see below). Nevertheless, the equations (1)–(4) may be used for determination of the ANC's if the following additional requirements [22]

$$\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)}) = f(E) \quad (5)$$

and

$$C_{l_f j_f}^2 = \frac{S_{l_f j_f}(E)}{\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})} = const \quad (6)$$

are fulfilled as a function of the free parameter $C_{l_f j_f}^{(sp)}$ for each energy E experimental point from the range $E_{min} \leq E \leq E_{max}$ and values of the function of $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ from (5).

One notes that the relations (5) and (6) can be considered as necessary and sufficient conditions of the fact that the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction are mainly peripheral. Hence, the relations (5) and (6) enables one to obtain the value $(C_{l_f j_f}^{exp})^2$ for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ using the experimental astrophysical S factors $S_{l_f j_f}^{exp}(E)$, precisely measured by authors of Refs. [6]–[10], instead of $S_{l_f j_f}(E)$, i.e.

$$(C_{l_f j_f}^{exp})^2 = \frac{S_{l_f j_f}^{exp}(E)}{\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})}. \quad (7)$$

Then, the value $(C_{l_f j_f}^{exp})^2$ can be used for extrapolation of the astrophysical S factor $S_{l_f j_f}(E)$ to the region of experimental inaccessible energies $0 \leq E < E_{min}$ by using the obtained value $(C_{l_f j_f}^{exp})^2$ in (1)–(4).

Values obtained in such a way for the $(C_{l_f j_f}^{exp})^2$ and $S_{l_f j_f}(E)$ at energies of $E < E_{min}$ can be considered as an "indirect determination" of the $(C_{l_f j_f}^{exp})^2$ for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ and of the astrophysical S factor for the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction at $E < E_{min}$, including $E = 0$. It should be noted that the expressions (1)–(4) allow one to determine both the absolute value of $(C_{l_f j_f}^{exp})^2$ for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ and that of the astrophysical S factor $S_{l_f j_f}(E)$ for the peripheral direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reactions in experimentally inaccessible energy regions by means of the analysis of the same precisely measured values of the experimental astrophysical S factors, $S_{l_f j_f}^{exp}(E)$ and $S_{34}^{exp}(E)$.

One notes that in the two-body potential model the ANC $C_{l_f j_f}$ is related to the NVC $G_{l_f j_f}$ for the virtual decay ${}^7\text{Be} \rightarrow \alpha + {}^3\text{He}$ by the equation [23]

$$G_{l_f j_f} = -i^{l_f + \eta_{7\text{Be}}} \frac{\sqrt{\pi}}{\mu} C_{l_f j_f}, \quad (8)$$

where the combinatorial factor taking into account the nucleon's identity is absorbed in $C_{l_f j_f}$ and its numerical value depends on a specific model used to describe wave functions of the ${}^3\text{He}$, α and ${}^7\text{Be}$ nuclei [24]. Hence, the proportionality factor in (8), which relates NVC's and ANC's, depends on the choice of nuclear model [24]. But, as it is noted in [24], the NVC $G_{l_f j_f}$ is a more fundamental quantity than the ANC $C_{l_f j_f}$ since the NVC is determined in a model-independent way as the residue of the partial S -matrix of the elastic $\alpha^3\text{He}$ -scattering at the pole $E = -\varepsilon_{\alpha^3\text{He}}$ ($\varepsilon_{\alpha^3\text{He}}$ is the binding energy of the bound $(\alpha + {}^3\text{He})$ state of ${}^7\text{Be}$) [23]. Therefore, values of the NVC could be obtained by the model-independent way from the analytic continuation in energy of $\alpha^3\text{He}$ -scattering amplitudes [24]. Besides, values of the NVC could also be obtained by the model-independent way from the method of extrapolation of the differential cross section for the ${}^3\text{He}({}^7\text{Be}, {}^3\text{He}){}^7\text{Be}$ reaction to the α -particle exchange singularity in the $\cos\theta$ plane [23] (θ is the scattering angle in the center of mass system). Values of the NVC obtained by these two quite different ways would allow one to get an additional information about correctness of the model approximations used for obtaining the relation (8). Therefore, it is also of interest to get an information about values of the NVC's from Eqs. (7) and (8).

2.2 The asymptotic normalization coefficients for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$

To determine the ANC values for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ the experimental astrophysical S factors, $S_{l_f j_f}^{exp}(E)$, for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction populating to the ground ($l_f=1$ and $j_f=3/2$) and first excited ($E^*=0.429 \text{ MeV}$; $J^\pi = 1/2^-$, $l_f=1$ and $j_f=1/2$) states are reanalyzed based on the relations (1)–(4), the conditions (5) and (6), and the relation (7). The recent experimental data measured by authors S.B. Nara Singh et al. [6], D. Bemmerer et al. [7], Gy. Gyüky et al. [8], F. Confortola et al. [9] and T.A.D. Brown et al. [10], which have apparently performed the most accurate direct measurement of the total astrophysical S factor for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction, are used for the analysis. These data cover the energy ranges $E=92.9$ – 168.9 keV [7–9], 420 – 951 keV [6] and 327 – 1235 keV [10] with absolute uncertainty not exceeding 5% and, consequently, the reaction under consideration is nonresonant. Also, one should note that the experimental astrophysical S factors for the the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction populating to the first and excited states of the ${}^7\text{Be}$ nucleus have been separated only for the energies of $E=92.9$, 105.6 and 147.7 keV in [9]. Whereas, in [10] the experimental astrophysical S factors have been separated for all experimental points of E from the aforesaid energy region by means of detecting the prompt γ ray (the prompt method) and by counting the ${}^7\text{Be}$ activity (the activation).

The Woods–Saxon potential split with a parity (l -dependence) for the spin-orbital term proposed by the authors of Refs. [31]–[33] is used here for the calculations of both bound state radial wave function $\varphi_{l_f j_f}(r)$ and scattering wave function $\psi_{l_i j_i}(r)$. It should be emphasized that the choice of this potential is based on the following considerations. Firstly, this potential form is justified from the microscopic point of view because it makes it possible to take into account the Pauli principle between nucleons in ${}^3\text{He}$ - and α -clusters in the $(\alpha + {}^3\text{He})$ bound state by means of inclusion of deeply bound states forbidden by the Pauli exclusion principle, i.e. without an explicit introduction of a repulsive core at small distance. The latter imitates the additional node (n) arising in the wave functions of $\alpha - {}^3\text{He}$ relative motion in ${}^7\text{Be}$. Secondly,

this potential describes well the phase shifts for $\alpha^3\text{He}$ -scattering in the wide energy range [32, 33] and reproduces the energies of low-lying states of the ^7Be nucleus [34].

The test of the peripheral character of the $^3\text{He}(\alpha, \gamma)^7\text{Be}$ reaction for the aforesaid energy range has been made by means of verifying the conditions (5) and (6), and by changing the geometric parameters (radius R and diffuseness a) of the adopted Woods–Saxon potential using the procedure of the depth adjusted to fit the binding energies, as it was done in Ref. [22]. According to Ref. [22], we vary R and a in the physically acceptable ranges (R in 1.62–1.98 fm and a in 0.63–0.77 fm) in respect to the standard values ($R=1.80$ fm and $a=0.70$ fm [32, 33]). Such a choice of the R and a parameters variation limit allows us to provide fulfillment of the conditions (5) and (6) in the aforesaid energy range within the experimental errors for the $S_{l_f j_f}^{exp}(E)$.

As an illustration, Fig.1 shows plots of the $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ dependence on the single-particle ANC, $C_{l_f j_f}^{(sp)}$ for $l_f=1$ and $j_f=3/2$ and $1/2$ only for the two values of energy E . The width of the band for these curves is the result of the weak "residual" (R, a)-dependence of the $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ on the parameters R and a (up to $\pm 2\%$) for the $C_{l_f j_f}^{(sp)} = C_{l_f j_f}^{(sp)}(R, a) = \text{const}$ [22, 30]. The same dependence is also observed at other energies. For example, for fig.1 plotted for $E=0.1056$ (0.1477) MeV overall uncertainty ($\Delta_{\mathcal{R}}$) of the function $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ in respect to the central values $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ corresponding to the central values of $C_{l_f j_f}^{(sp)} (= C_{l_f j_f}^{(sp)}(1.80, 0.70))$ comes to $\Delta_{\mathcal{R}} = \pm 4.5(\pm 4.5)\%$ for the ground state of ^7Be and $\Delta_{\mathcal{R}} = \pm 3.4(\pm 2.9)\%$ for the excited state of ^7Be . As it is seen from here that the $^3\text{He}(\alpha, \gamma)^7\text{Be}(0.429 \text{ MeV})$ reaction is slightly more peripheral than the $^3\text{He}(\alpha, \gamma)^7\text{Be}(\text{g.s.})$ reaction since the binding energy for $^7\text{Be}(0.429 \text{ MeV})$ is less than that for $^7\text{Be}(\text{g.s.})$. The similar dependence of $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ on the $C_{l_f j_f}^{(sp)}$ values is observed for other energies and the value of $\Delta_{\mathcal{R}}$ is no more than $\pm 5.0\%$ in respect to the central values $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ within the interval of $3.205 \leq C_{13/2}^{(sp)} \leq 4.397 \text{ fm}^{-1/2}$ ($2.788 \leq C_{11/2}^{(sp)} \leq 3.763 \text{ fm}^{-1/2}$) for the $^3\text{He}(\alpha, \gamma)^7\text{Be}(\text{g.s.})$ ($^3\text{He}(\alpha, \gamma)^7\text{Be}(0.429 \text{ MeV})$) reaction. These ranges of change for $C_{l_f j_f}^{(sp)}$ correspond to the parameters of the adopted Woods–Saxon potential R ranging from 1.62–1.98 fm and a in the range of 0.63–0.77 fm. It follows from here that the condition (5) is satisfied for the considered reaction within the uncertainties not exceeding the experimental errors of $S_{l_f j_f}^{exp}(E)$.

We also calculated the $\alpha^3\text{He}$ -elastic scattering phase shifts by variation of the parameters R and a in the same range for the adopted Woods–Saxon potential. The results of the calculations corresponding to s - and p -waves are presented in Fig.2 in which the width of the bands corresponds to a change of phase shifts values with respect to variation of values of the R and a parameters. As it is seen from Fig.2, the experimental phase shifts [35] are well reproduced within uncertainty of about $\pm 5\%$.

This circumstance allows us to test the condition (6), which is no less essential for the peripheral character of these reactions, at the energies of $E=92.9, 105.6$ and 147.7 keV for which the $^3\text{He}(\alpha, \gamma)^7\text{Be}(\text{g.s.})$ and $^3\text{He}(\alpha, \gamma)^7\text{Be}(0.429 \text{ MeV})$ astrophysical S factors were separately measured in [9]. As an illustration, for the same energies E as in Fig.1 we present in Fig.3 (the upper panels) the results of $C_{l_f j_f}^2$ -value calculation given by Eq.(6) ($(l_f j_f)=(1 \ 3/2)$ and

(1 1/2)) in which instead of the $S_{l_{fj_f}}(E)$ the experimental S factors for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction populating to the ground and first excited states of ${}^7\text{Be}$ were taken. It should be noted that the same dependence occurs for other considered energies. The calculation shows the obtained $C_{l_{fj_f}}^2$ values also weakly (up to 5.0 %) depend on the $C_{l_{fj_f}}^{(sp)}$ -value. However, the values of the spectroscopic factors $Z_{1\ 3/2}$ and $Z_{1\ 1/2}$ corresponding to the $(\alpha + {}^3\text{He})$ -configuration for ${}^7\text{Be}(\text{g.s.})$ and ${}^7\text{Be}(0.429\text{ keV})$, respectively, change strongly about 1.7 times (see, the lower panels in Fig.3).

Thus, the calculation shows that the uncertainty in $\mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$ and $C_{l_{fj_f}}^2$ is up to $\pm 5.0\%$ relative to the central values of $\mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$ and $C_{l_{fj_f}}^2$, obtained for the standard values of $R = 1.80\text{ fm}$ and $a = 0.70\text{ fm}$, for the (R, a) -pair varying in the above mentioned intervals for R and a , while the uncertainty in the $Z_{l_{fj_f}}$ is about $\pm 35\%$.

For different energies E we also estimate a relative contribution of the nuclear interior ($r \leq r_N$) to the astrophysical S factors for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction populating to the ground and first excited states in dependence on the variation $C_{l_{fj_f}}^{(sp)}$ (or R and a) introducing the cutoff radius r_{cut} ($r_{cut} \approx r_N$) in the lower limit of integration of the radial integral (10) of Ref.[22] entering in the amplitude of the reaction under consideration. With this aim one considers the ratio $\Delta(E, C_{l_{fj_f}}^{(sp)}; r_{cut}) = | \mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)}) - \tilde{\mathcal{R}}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)}; r_{cut}) | / \mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$, where $\tilde{\mathcal{R}}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)}; r_{cut})$ is given by Eqs.(10) and (13) of Ref.[22] but in the radial integral (10) of Ref.[22] the integration over r is performed in the interval $r_{cut} \leq r \leq \infty$, i.e. $\tilde{\mathcal{R}}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)}; 0) = \mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$. The $\mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$ and $\tilde{\mathcal{R}}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)}; r_{cut})$ functions were calculated for different values of the single-particle ANC $C_{l_{fj_f}}^{(sp)}$ (or the parameters R and a). A value of the cutoff radius is taken as in Ref. [36], that is $r_{cut} = r_N = 1.36(4^{1/3} + 3^{1/3}) = 4.12\text{ fm}$, as well as $r_{cut} = 4.00\text{ fm}$ and 4.25 fm . The calculation of $\Delta(E, C_{l_{fj_f}}^{(sp)}; r_{cut})$ performed at different energies shows that the quantities of $\Delta(E, C_{l_{fj_f}}^{(sp)}; r_{cut})$ change from 5.4% up to 15.5% under variation of $C_{l_{fj_f}}^{(sp)}$ and r_{cut} . The calculation shows that the contribution of the nuclear interior ($r \leq r_N$) to the astrophysical S factors calculated for different sets of geometric parameters R and a of the Woods–Saxon potential, and values of r_{cut} does not exceed about 15.5% and this small quantity is due mainly to oscillations observed in the integrand of the radial integral (10) of Ref.[22]. As an illustration of this fact, in Fig.4 we show a dependence of the integrand of the radial integral (10) of Ref.[22] on geometric parameters R and a of the Woods–Saxon potential and values of r_{cut} for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ reaction at different energies. As one can see from Fig.4 the integrand of the radial integral changes with the variation of the geometric parameters R and a , which is associated with changes of the calculated bound $(\alpha + {}^3\text{He})$ state wave function and the calculated $\alpha^3\text{He}$ -scattering wave function, and these wave functions indeed reached simultaneously their asymptotic form for $r \gtrsim 5.0\text{ fm}$. Such a change leads to calculated $\tilde{S}_{l_{fj_f}}(E)$ that vary by 1.75 times over the energy region $92.9 \leq E \leq 1200\text{ keV}$, while the calculated values of the function $\mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$ change by only $\pm 5\%$ with respect to the value of $\mathcal{R}_{l_{fj_f}}(E, C_{l_{fj_f}}^{(sp)})$ corresponding to the standard values of $R = 1.80\text{ fm}$ and $a = 0.70\text{ fm}$. Besides, it is seen from Fig.4 that the behavior of the integrand in the radial integral (10) of

Ref.[22] over a wide energy range provides a strong suppression of the contribution only from the part of the nuclear interior with $0 \leq r \lesssim 2.0$ fm to the integral (10) of Ref.[22]. However, a noticeable change of the integrand of the aforesaid integral is observed with the variation of the parameters R and a in the range $2.0 \lesssim r \lesssim 6.0$ fm. At this the contributions from the regions with $2.0 \lesssim r \lesssim 4.0$ fm and $4.0 \lesssim r \lesssim 6.0$ fm can compensate one another because of the presence of the noticeable oscillation in the integrand. Moreover, as it is seen from Fig.4, the lower and upper limits of this range depend on the parameters R and a , and the choice of the cutoff radius r_{cut} also becomes ambiguous since a fitted value of r_{cut} , which in turn becomes dependent on the parameters R and a . The similar situation occurs for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ reaction. Such behavior of the integrand in the radial integral allows one to understand one of the main reason why the aforesaid phenomenological values of the spectroscopic factors (the ANC's) become noticeably (weekly) dependent from a choice of values of the free parameters R and a (or the free parameter $C_{1j_f}^{(sp)}$). Therefore, the parametrization of the astrophysical S factors in terms of the spectroscopic factors cannot be applied for the ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be} + \gamma$ reaction and the expressions (1)–(4) should be used for the analysis to reduce the above mentioned ambiguities to minimum. In this connection one would like to note the following. In paper [19], as it is mentioned above, the calculation of the astrophysical S factors for the reactions under consideration has been carried out using the expression (1) but introducing the cutoff radius r_{cut} in the lower limit of integration in the radial integral (10) of Ref.[22] and replacing the bound state wave function $\varphi_{l_f j_f}(r)$ with its asymptotic form starting from $r = r_{cut}$. At this the best fitting of the calculated $S_{l_f j_f}(E)$ to the experimental ones [25] was reached when the cutoff radius was $r_{cut}=4.0$ fm. It is seen from here that in paper [19] the contribution of the nuclear interior to the calculated astrophysical S factors was indeed underestimated, since contribution of the nuclear interior $0 < r \leq 4.0$ fm to the calculated astrophysical S factors, which is up to about 14%, has not been taken into account. Here, firstly, the contribution of the nuclear interior ($r \leq r_N$) to the calculated astrophysical S factors is taken into account in a correct way by means of the appropriate choice of the adopted potential both for the initial state and for final state of the reactions under consideration. Secondly, the problem of the ambiguity connected with the strong (R, a) -dependence of the calculated astrophysical S factors is removed by inclusion of the information about ANC. The latter reduces this ambiguity to minimum. At last, in the present work the more precise experimental data for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factors [6–10] than those in [19] are used for the analysis. Here, using by the case, one notes that there is a misprint in the line 36 upper of section 3 of [19]. There the phrase "no more than 1% to" must be written as "no more than 10% to".

Thus, the scrupulous analysis performed here quantitatively shows that the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction within the considered energy ranges is peripheral and the dominant contribution into the amplitude gives regions well outside the range of the internuclear $\alpha{}^3\text{He}$ interactions. Hence, a use of parametrization in terms of the ANC's is adequate to the physics of the reaction under consideration since ANC for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}$ determines the probability of finding the $(\alpha + {}^3\text{He})$ configuration in ${}^7\text{Be}$ at distances between fragments α and ${}^3\text{He}$ that larger than the range of their nuclear interaction.

For each energy E experimental point ($E=92.9, 105.6$ and 147.7 keV) the values of the ANC's are obtained for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(g.s.)$ and $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ by using the

corresponding experimental astrophysical S factor ($S_{1\,3/2}^{exp}(E)$ and $S_{1\,1/2}^{exp}(E)$, (the activation)) [7, 9] in the ratio of the r.h.s. of the relation (1) instead of the $S_{l\,j\,j_f}(E)$ and the central values of $\mathcal{R}_{l\,j\,j_f}(E, C_{l\,j\,j_f}^{(sp)})$ corresponding to the adopted values of the parameters R and a . The results of the ANC's, $(C_{1\,3/2}^{exp})^2$ and $(C_{1\,1/2}^{exp})^2$, for these three energy E experimental points are displayed in Fig.5 (filled circle symbols) and the second and third columns of Table 1. The uncertainties pointed in this figure correspond to those found from (1) (averaged square errors (a.s.e.)), which include the total experimental errors in the corresponding experimental astrophysical S factor and the aforesaid uncertainty in the $\mathcal{R}_{l\,j\,j_f}(E, C_{l\,j\,j_f}^{(sp)})$. One should note that the same results for the ANC's are obtained when $S_{34}^{exp}(E)$ ($S_{1\,3/2}^{exp}(E)$ and $R^{exp}(E)$) [7, 9] are used in Eq.(2) (in Eq.(3) and (4)) instead of $S_{34}(E)$ ($S_{1\,3/2}(E)$ and $R(E)$). Then in Eq.(3), inserting the averaged means of λ_C ($\lambda_C=0.666$), obtained from the three data, and replacing of the $S_{34}(E)$ in the l.h.s. of Eq.(3) with $S_{34}^{exp}(E)$ for the others, the two experimental points of energy E ($E=126.5$ and 168.9 keV) from [7, 8], four one E ($E=420.0, 506.0, 615.0$ and 951.0 keV) from [6] and the three one E ($E=93.3, 106.1$ and 170.1 keV) from [9] can also determine values of ANC's, $C_{1\,3/2}^2$ and $C_{1\,1/2}^2$. The results of the ANC's ($C_{1\,j\,j_f}^{exp})^2$ for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be(g.s.)}$ and $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429\text{ MeV})$ are displayed in Fig.5 (the opened cycle and triangle symbols) and the second and third columns of Table 1. The same way the values of the ANC's are obtained by using the separated experimental astrophysical S factors ($S_{1\,3/2}^{exp}$ and $S_{1\,1/2}^{exp}$) [10]. The results for these ANC's are also presented in Figs.5 both for the activation (filled star symbols) and for the prompt method (filled square symbols). Also, in Fig.5 and Table 2 the weighted means of the ANC's-values and their uncertainties (the solid line and the band width, respectively), derived both separately from each experimental data and from all of the experimental points, are presented. As it is seen from Fig.5 and Tables 1 and 2 the values of the ANC's, obtained from the analysis of the experimental astrophysical S factors measured by the authors of Refs.[6–10] in different intervals of energy E , agree rather well with each other. It should be noted that the ratio in the r.h.s. of the relation (7) does not practically depend on the energy E , although absolute values of the corresponding experimental astrophysical S factors for the reactions under consideration depend noticeably on the energy and change by up to about 1.7 times when E changes from 92.6 keV to 1200 keV. This fact allows us to conclude that the energy dependence of the experimental astrophysical S factors [6]–[10] is well determined by the calculated function $\mathcal{R}_{l\,j\,j_f}(E, C_{l\,j\,j_f}^{(sp)})$ and $\mathcal{R}_{13/2}(E, C_{13/2}^{(sp)}) + \lambda_C \mathcal{R}_{11/2}(E, C_{11/2}^{(sp)})$. Hence, the experimental astrophysical S factors measured by the authors of Refs.[6]–[9] can be used as an independent source of reliable information about the ANC's for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be(g.s.)}$ and $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429\text{ MeV})$.

As it is seen from the first and second (third and fourth) lines of Table 2 the weighted means of the ANC's-values for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be(g.s.)}$ and $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429\text{ MeV})$ obtained by the analysis performed separately for the activation and the prompt method of the experimental data from the works [6]–[9] and the work [10] are in a good agreement with one another and their uncertainties do not exceed about 3% for Refs.[6]–[9] and 6% for Ref.[10]. While the weighted means of the ANC-values found by using separately the experimental data of works Refs.[6]–[9] and the work Ref.[10] noticeably differ from one another (up to 1.13 times for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be(g.s.)}$ and up to 1.12 times for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429\text{ MeV})$). The main reason of the

latter is in the observed difference in absolute values of the experimental astrophysical S factors measured by authors of works Refs [6]–[9] and of the work Ref. [10] (see Fig.6). Nevertheless, the weighted means of the ANC-values recommended by us for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ obtained from all of the experimental data are equal to $(C_{1\,3/2}^{\text{exp}})^2 = 23.2 \pm 1.4 \text{ fm}^{-1}$ and $(C_{1\,1/2}^{\text{exp}})^2 = 15.7 \pm 1.0 \text{ fm}^{-1}$ ($C_{1\,3/2}^{\text{exp}} = 4.82 \pm 0.15 \text{ fm}^{-1/2}$ and $C_{1\,1/2}^{\text{exp}} = 3.97 \pm 0.13 \text{ fm}^{-1/2}$), respectively. The overall uncertainties for the ANC's recommended by us for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ are about 6.0%, which are a bit more than the aforesaid uncertainties obtained from the separate analysis of the experimental astrophysical S factors measured by authors of Refs.[6]–[9] and Ref.[10]. This is connected with the aforesaid discrepancy between the ANC-values presented in the first and second lines as well as the fifth and sixth lines of Table 2. As it is seen from Table 2, the weighted means of the ANC-values for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ recommended by us are from 1.2σ to 1.4σ level larger (from 0.4σ to 0.6σ level lower) and from 0.9σ to 1.1σ level larger (from 0.4σ to 0.6σ level lower) than the weighted means of the ANC-values deduced by us from the separate analysis of the experimental astrophysical S factors from works [6]–[9] (work [10]), respectively. The corresponding values of the NVC's are $|G_{1\,3/2}|^2 = 1.10 \pm 0.07 \text{ fm}$ and $|G_{1\,1/2}|^2 = 0.742 \pm 0.047 \text{ fm}$. These results for the ANC's and NVC's recommended by us are the central ones of this paper.

One should note that the values of $C_{1\,3/2}$ and $C_{1\,1/2}$ should not be equal, in contrast with the assumption made in Ref.[11]. As noted earlier in paper [19], the values ANC's (NVC's) $C_{1\,3/2}^2 = 18.2 \text{ fm}^{-1}$ and $C_{1\,1/2}^2 = 15.0 \text{ fm}^{-1}$ ($|G_{1\,3/2}|^2 = 0.860 \text{ fm}$ and $|G_{1\,1/2}|^2 = 0.709 \text{ fm}$) were obtained from the experimental data analysis [25], which has considerable spread. Also, as mentioned above, in paper [19] the contribution of the nuclear interior ($r < 4.0 \text{ fm}$) to the calculated astrophysical S factors was not included. It is seen that taking into account the contribution of the nuclear interior and use the experimental data more accurate than those in Ref.[19] one can noticeably influence the extracted values of the ANC's. A comparison of the present result and that obtained in paper [19] shows that the underestimation of the contribution both of the nuclear interior and of the nuclear exterior indeed occurs in [19] since the present value of ANC $C_{1\,3/2}^2$ obtained from the analysis of the more accurate experimental astrophysical S factor [6-10] is larger than that obtained in [19].

The resulting ANC (NVC) value for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(\text{g.s.})$ obtained by us is in good agreement with the value $C_{1\,3/2}^2 = 25.3 \text{ fm}^{-1}$ ($|G_{1\,3/2}|^2 = 1.20 \text{ fm}$) and that for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ differs noticeably from the value $C_{1\,1/2}^2 = 22.0 \text{ fm}^{-1}$ ($|G_{1\,1/2}|^2 = 1.04 \text{ fm}$), which were obtained in [37] within the $(\alpha + {}^3\text{He})$ -channel resonating-group method. Also, the results recommended by us for these ANC's differ noticeably from the values $C_{1\,3/2}^2 = 12.6 \pm 1.1 \text{ fm}^{-1}$ and $C_{1\,1/2}^2 = 8.41 \pm 0.58 \text{ fm}^{-1}$ ($C_{1\,3/2} = 3.55 \pm 0.15 \text{ fm}^{-1/2}$, $C_{1\,1/2} = 2.90 \pm 0.10 \text{ fm}^{-1/2}$, $|G_{1\,3/2}|^2 = 0.596 \pm 0.052 \text{ fm}$ and $|G_{1\,1/2}|^2 = 0.397 \pm 0.030 \text{ fm}$) [15] as well as those $C_{1\,3/2}^2 = C_{1\,1/2}^2 = 14.4 \text{ fm}^{-1}$ ($C_{1\,3/2} = C_{1\,1/2} = 3.79 \text{ fm}^{-1/2}$ and $|G_{1\,3/2}|^2 = |G_{1\,1/2}|^2 = 0.680 \text{ fm}$) [11]. In this connection one would like to draw attention to the following. The bound state wave functions and the initial state wave functions in [15] were computed with different potentials and, so, these calculations were not self-consistent. Besides, the values of the binding energies for the bound states of ${}^7\text{Be}$ calculated in Ref.[15] differ from the experimental ones. Therefore, the calculated value of

the binding energy for the bound state of ${}^7\text{Be}(\text{g.s.})$ in the $(\alpha + {}^3\text{He})$ -channel (4.73 MeV, see Table I in Ref.[15]) is also not in agreement with the experimental one (1.59 MeV). Since the ANC's (or NVC's) for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ are sensitive to the form of the NN potential, it is desirable, firstly, to calculate the wave functions of the bound state using other forms of the NN potential, and, secondly, in order to guarantee the self-consistency, the same forms of the NN potential should be used for such calculation of the initial wave functions. Besides, one would also like to note the recent result of Ref.[11] obtained for $C_{1\,3/2}$ and $C_{1\,1/2}$ from the analysis of the experimental ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factors performed within the R -matrix method, where the contribution from the internal part of the amplitude was simulated by the background for a single pole. But there to reduce the number of free parameters the assumption about equality of the ANC's ($C_{1\,3/2} = C_{1\,1/2}$) was used, and the best fitting of the data was reached at $C_{1\,3/2}=3.79\text{ fm}^{-1/2}$ and the channel radius $r_c=3.0\text{ fm}$. It follows from here that in reality the values of the ANC's, $C_{1\,3/2}$ and $C_{1\,1/2}$, should not be equal. Moreover, the calculation shows that the asymptotic behavior of the bound $(\alpha + {}^3\text{He})$ state and $\alpha{}^3\text{He}$ -scattering wave functions is reached, as it was mentioned above, simultaneously only at $r_c \gtrsim 5.0\text{ fm}$ and, so, at $r_c \geq 3.0\text{ fm}$ their substitution for these wave functions in the external part of the amplitude in Ref.[11] is not correct.

2.3 α -particle spectroscopic factors for the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair

The "indirectly determined" values of the ANC's for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ obtained in the present work and those for $\alpha + t \rightarrow {}^7\text{Li}$ deduced in Ref.[22] can be used for obtaining information on the ratio $R_{Z;j_f} = Z_{1j_f}({}^7\text{Be})/Z_{1j_f}({}^7\text{Li})$ for the virtual α decays of the bound mirror (${}^7\text{Li}{}^7\text{Be}$)-pair, where $Z_{1j_f}({}^7\text{Be})(Z_{1j_f}({}^7\text{Li}))$ is the spectroscopic factor for ${}^7\text{Be}$ (${}^7\text{Li}$) in the $(\alpha + {}^3\text{He})((\alpha + t))$ -configuration. For this aim, from $C_{1j_f}(\text{B}) = Z_{1j_f}^{1/2}(\text{B})C_{1j_f}^{(sp)}(\text{B})$ ($\text{B} = {}^7\text{Li}$ and ${}^7\text{Be}$) we form the relation

$$R_{Z;j_f} = \frac{R_{C;j_f}}{R_{C^{(sp)};j_f}}, \quad (9)$$

where $R_{C;j_f} = \left(C_{1j_f}({}^7\text{Be})/C_{1j_f}({}^7\text{Li})\right)^2$ ($R_{C^{(sp)};j_f} = \left(C_{1j_f}^{(sp)}({}^7\text{Be})/C_{1j_f}^{(sp)}({}^7\text{Li})\right)^2$) is the ratio of squares of the ANC's (single-particle ANC's) for the bound mirror (${}^7\text{Li}{}^7\text{Be}$)-pair and $j_f=3/2(1/2)$ for the ground (first excited) state of the mirror nuclei. Besides, it should be noted that the relation (9) allows one to verify a validity of the approximation ($R_{C;j_f} \approx R_{C^{(sp)};j_f}$, i.e., $R_{Z;j_f} \approx 1$) used in Refs.[38] for the mirror (${}^7\text{Li}{}^7\text{Be}$) conjugated α decays.

For the bound and first excited state of the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair the values of $C_{1j_f}^{(sp)}({}^7\text{Be})$ and $C_{1j_f}^{(sp)}({}^7\text{Li})$ change by the factor of 1.3 under the variation of the geometric parameters (R and a) of the adopted Woods-Saxon potential [32, 33] within the aforesaid ranges, while the ratios $R_{C^{(sp)};3/2}$ and $R_{C^{(sp)};1/2}$ for the bound and first excited states of the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair change by only about 1.5% and 6%, respectively. It is seen that the ratios do not depend practically from variation of the free parameters R and a . Therefore, these ratios can be determined in model-independent way by using the values of the single-particle ANC's for the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair obtained in the present work and deduced in Ref.[22]. The ratios

are equal to $R_{C^{(sp)};3/2}=1.37\pm 0.02$ and $R_{C^{(sp)};1/2}=1.40\pm 0.09$, which are in good agreement with those calculated in [38] within the microscopic cluster and two-body potential models (see Table I there). The ratios for the ANC's are $R_{C;3/2}=1.85\pm 0.11$ and $R_{C;1/2}=1.75\pm 0.10$. From (9) the values of the ratio $R_{Z;j_f}$ are equal to $R_{Z;3/2}=1.35\pm 0.08$ and $R_{Z;1/2}=1.25\pm 0.11$ for the ground and the first excited states, respectively. These values differ noticeably from those of $R_{Z;3/2}=0.995\pm 0.005$ and $R_{Z;1/2}=0.990$ calculated in Ref.[38] within the microscopic cluster model. One notes that the values of $R_{Z;j_f}$ calculated in [38] are sensitive to the model assumptions (the choice of the oscillation radius b and the form of the effective NN potential) and such model dependence may actually influence the mirror symmetry for the α -particle spectroscopic factors. The mirror symmetry breakup for the α -particle spectroscopic factors can also be signalled by the results for the ratio of $S_{34}(^7\text{Be})/S_{34}(^7\text{Li})$ at zero energies for the mirror ($^7\text{Li}^7\text{Be}$)-pair obtained in [12] within the resonating-group method by using the seven forms for the effective NN potential. As it is seen from Ref.[12], this ratio is sensitive to a form of the effective NN potential used and changes from 1.0 to 1.18 times in a dependence from the effective NN potential used. One of the possible reasons of the sensitivity observed in [12] can apparently be associated with a sensitivity of the ratio $R_{Z;j_f}$ to a form of the effective NN potential used. In a contrast of such model dependence observed in [12, 38], the problem of the ambiguity connected with the model (R, a) -dependence for the values of the ratios $R_{Z;j_j}$ found by us from Eq.(9) is reduced to minimum within the experimental uncertainty.

It is seen from here that the empirical values of $R_{Z;j_f}$ noticeably exceed unity both for the ground state and for the first excited state of the mirror ($^7\text{Li}^7\text{Be}$)-pair. This result for $R_{Z;j_f}$ is not accidental and can be explained qualitatively by the following consideration. The fact is that the spectroscopic factor $Z_{1j_f}(^7\text{Li})$ (or $Z_{1j_f}(^7\text{Be})$) is determined as a norm of the radial overlap function of the bound state wave functions of the t , α and ^7Li (or ^3He , α and ^7Be) nuclei and is given by Eqs.(100) and (101) from Ref. [23]. The interval of integration ($0 \leq r < \infty$) in Eq. (101) can be divided in two parts. In the first integral denoted by $Z_{1j_f}^{(1)}(^7\text{Li})$ for ^7Li and $Z_{1j_f}^{(1)}(^7\text{Be})$ for ^7Be , the integration over r covers the region $0 \leq r \leq r_N$ (the internal region), where nuclear (αt or $\alpha^3\text{He}$) interactions are dominate over the Coulomb interactions. In the second integral

$$Z_{1j_f}^{(2)}(B) = C_{1j_f}^2(B) \int_{r_N}^{\infty} dr W_{\eta_B;3/2}^2(2\kappa_{\alpha\alpha}r), \quad (10)$$

where in the asymptotic region the radial overlap function entering the integrand is replaced by the appropriate Whittaker function (see, for example, Eq.(108) of Ref.[23]), interaction between a and α -particle ($a = t$ for $B = ^7\text{Li}$ or $a = ^3\text{He}$ for $B = ^7\text{Be}$) is governed by the Coulomb forces only (the external region). In (10), $\kappa_{\alpha\alpha} = \sqrt{2\mu_{\alpha\alpha}\varepsilon_{\alpha\alpha}}$ and $W_{\eta_B;3/2}(x)$ is the Whittaker function. One notes that the magnitudes $Z_{1j_f}^{(1)}(^7\text{Li})$ ($Z_{1j_f}^{(1)}(^7\text{Be})$) and $Z_{1j_f}^{(2)}(^7\text{Li})$ ($Z_{1j_f}^{(2)}(^7\text{Be})$) define the probability of finding t (or ^3He) in the $(\alpha + t)$ configuration (or the $(\alpha + ^3\text{He})$ configuration) at distances of $r \leq r_N$ and of $r > r_N$, respectively. Obviously $Z_{1j_f}(^7\text{Li}) = Z_{1j_f}^{(1)}(^7\text{Li}) + Z_{1j_f}^{(2)}(^7\text{Li})$ and $Z_{1j_f}(^7\text{Be}) = Z_{1j_f}^{(1)}(^7\text{Be}) + Z_{1j_f}^{(2)}(^7\text{Be})$.

An information about values of $Z_{1j_f}^{(2)}(^7\text{Li})$ and $Z_{1j_f}^{(2)}(^7\text{Be})$ can be obtained from (10) by using the values of the ANC's for $\alpha + t \rightarrow ^7\text{Li}$ and $\alpha + ^3\text{He} \rightarrow ^7\text{Be}$ recommended in [22] and in the

present work, respectively. For example, for $r_N \approx 4.0$ fm (the surface regions for the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair) the calculation shows that the ratio $R_{Z;j_f}^{(2)} = Z_{1j_f}^{(2)}({}^7\text{Be})/Z_{1j_f}^{(2)}({}^7\text{Li})$ is equal to 1.42 (1.29) for the ground (excited) states of the ${}^7\text{Li}$ and ${}^7\text{Be}$ nuclei, i.e. the ratio $R_{Z;j_f}^{(2)} > 1$. Owing to the principle of equivalency of nuclear interactions between nucleons of the (αt) -pair in the nucleus ${}^7\text{Li}$ and $(\alpha {}^3\text{He})$ -pair in the nucleus ${}^7\text{Be}$ [38], the values of $Z_{1j_f}^{(1)}({}^7\text{Li})$ and $Z_{1j_f}^{(1)}({}^7\text{Be})$ should not differ appreciably. If one suggests that $R_{Z;j_f}^{(1)} \approx 1$, then the ratio $R_{Z;j_f} > 1$.

2.4 Astrophysical S factor for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction at solar energies

The equation (1) and the weighted means of the ANC's obtained for the ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ can be used for extrapolating the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factor for capture to the ground and first excited states as well as the total astrophysical S factor at solar energies ($E \leq 25$ keV). At first, we tested again the fulfilment of the condition (5) in the same way as it is done above for $E \geq 90$ keV. Similar results plotted in Fig.1 are also observed for the $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ function dependence on the single-particle ANC, $C_{l_f j_f}^{(sp)}$, at energies of $E < 90$ keV.

The experimental and calculated astrophysical S factors for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$, ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.} + 0.429 \text{ MeV})$ reactions are presented in Table 1 and displayed in Fig.6a, b and c, respectively. The symbols in these figures are data of all experiments [6]–[10] and the results of the present work. In Figs.6a and b, the opened diamond and triangle symbols show our results for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ reactions (see Table 1 also), which are obtained from the analysis of the total experimental astrophysical S factors of [7-9] and [6] by using the corresponding values of the ANC's for each energy E experimental point presented in Table 1. These results are the second ones of the present work. In Figs.6a and b the experimental data plotted by the filled circle symbols (filled star and square symbols) are taken from [9] (from [10]). The opened circle symbols in these figures are the results of the extrapolation recommended in the present work in which each of the quoted uncertainties is the uncertainty associated with that for the ANC's adopted. The solid lines present our calculations performed with the standard values of geometric parameters $R=1.80$ fm and $a=0.70$ fm both for the bound $(\alpha + {}^3\text{He})$ state and for $\alpha {}^3\text{He}$ -scattering state. In Fig.6(c), the symbols are data of all experiments [6]–[10] and the present work (the open cycle symbols) as well as the dashed line is the result of Ref.[16]. Everywhere the width of each of the band is the weighted uncertainty, which is associated with the uncertainties in the "indirectly determined" values of ANC's. As it is seen from these figures, the equations (1), (2) and (7) allow one to perform a correct extrapolation of the corresponding astrophysical S factors at solar energies when the corresponding ANC-values are known. But, the noticeable underestimation between the result of calculation performed in Ref[16] in respect to the experimental data (the dashed line in Fig.6c) occurs.

The weighted means of the total astrophysical S factor $S_{34}(E)$ at solar energies ($E=0$ and 25 keV b) are presented in Table 2. As it is seen from Table 2 the weighted means of $S_{34}(0)$, deduced by us separately from each the activation and the prompt method of the

experimental data from works [6]–[9] and work [10], agree well within their uncertainties with those recommended in [9, 10]. The weighted mean of $S_{34}(0)$ obtained from those presented in the first and second (the fifth and sixth) lines of Table 2 is equal 0.628 ± 0.018 keV b (0.567 ± 0.004 keV b) and they also agree reasonably well within the corresponding uncertainty with those recommended in [10] and [9]. But, these weighted means of $S_{34}(0)$ noticeably differ from one another (about 11%). The main reason of this distinction is in the observed difference in magnitudes of the ANC's presented in Table 2, which is in turn associated with the observed discrepancy between the experimental astrophysical S factors measured by authors of Refs [6]–[9] and of Ref.[10], as it is mentioned above. Nevertheless, the weighted mean of $S_{34}(0) = 0.610 \pm 0.037$ keV b, recommended by us and deduced from all the weighted means of $S_{34}(0)$ presented in Table 2, is about 0.4σ (1.4σ) level larger than those recommended in [10] ([9]), respectively. But, the overall uncertainty for $S_{34}(0)$ recommended by us is about 6%, whereas the uncertainty recommended by authors of works [9] (work [10]) is about 3% (about 4%). The main reason of this noticeable distinction is associated with the overall uncertainty (about 6%) in the weighted mean of the ANC's for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ (see the last line of Table 2), which is obtained from the analysis of all the experimental data [6–10]. Comparison of our result with that obtained in [19, 15] shows that a noticeable discrepancy between the present results and those of Refs.[19, 15] occurs. This circumstance is apparently connected with the underestimated value of $C_{1\,3/2}^2$ and $C_{1\,1/2}^2$ (or $|G_{1\,3/2}|^2$ and $|G_{1\,1/2}|^2$) obtained in Ref.[19, 15] in respect to our result. Besides, one should note that our result differs also from that recommended in Refs.[11, 6] ($S_{34}(0)=0.51 \pm 0.04$ keV b [11] and 0.53 keV b [6]). This difference is a result of the underestimation of the contribution from the external part in the amplitude admitted in [11]. One notes once more that in [6] the value of $S_{34}(0)$ has been obtained by the artificial renormalization of the result of [11].

Besides, the result of the present work is noticeably larger than the result of $S_{34}(0)=0.516$ keV [16] obtained within the standard two-body ($\alpha + {}^3\text{He}$) potential by using $\alpha {}^3\text{He}$ potential deduced by a double-folding procedure. One of the possible reason of this discrepancy can be apparently associated with the assumption admitted in [16] that a value of the ratio $R_{Z;j_f}$ for the bound mirror (${}^7\text{Li}{}^7\text{Be}$)-pair is taken equal to unity ($Z_{13/2}({}^7\text{Be}) = Z_{11/2}({}^7\text{Be})=1.17$ [39]), which in turn results in the observed systematical underestimation of the calculated $S(E)$ in respect to the experimental S factors (see the dashed line in Fig.6c). But, as shown in Subsection 2.3 the values of $R_{Z;j_f}$ for the ground and first excited states of the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair are noticeably larger unity. Therefore, the underestimated values of $Z_{13/2}({}^7\text{Be})$ and $Z_{11/2}({}^7\text{Be})$ used in [16] also result in the underestimated value of $S_{34}(0)$ for the direct capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction. Perhaps, the assumption about equal values of the spectroscopic factor ($Z_{13/2}({}^7\text{Li}) = Z_{11/2}({}^7\text{Li})=1.17$ [16, 39]) is correct only for the spectroscopic factors $Z_{1j}({}^7\text{Li})$ since the value of $S_{34}(0)$ obtained in [22] and [16] for the direct capture $t(\alpha, \gamma){}^7\text{Li}$ reaction agree excellently with each other. One notes that in Ref.[22] the analysis of the the $t(\alpha, \gamma){}^7\text{Li}$ reaction has also been performed within MTBPA and the ANC for $\alpha + t \rightarrow {}^7\text{Li}(\text{g.s.})$ presented in [40], which has been deduced there from the results of Ref.[16], also is in good agreement with that recommended by authors of Ref.[22].

Nevertheless, we observe that the value of $S_{34}(0)=0.56$ keV b [13] obtained within the microscopical ($\alpha + {}^3\text{He}$)-cluster approach is in agreement with our result about 1.4σ level.

Besides, our result is also in excellent agreement with that of $S_{34}(0)=0.609$ keV b [12] and $S_{34}(0)=0.621$ keV b [37] obtained within the $(\alpha + {}^3\text{He})$ -channel of version of the resonating-group method by using the modified Wildermuth-Tang (MWT) and the near-Serber exchange mixture forms for the effective NN potential, respectively. It follows from here that the mutual agreement between the results obtained in the present work and works of [12, 13, 37], which is based on the common approximation about the cluster $(\alpha + {}^3\text{He})$ structure of the ${}^7\text{Be}$, allows one to draw a conclusion about the dominant contribution of the $(\alpha + {}^3\text{He})$ clusterization to the low-energy ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ cross section both in the absolute normalization and in the energy dependence [6–10]. Therefore, single-channel $(\alpha + {}^3\text{He})$ approximation for ${}^7\text{Be}$ [12, 13] is quite appropriate for this reaction in the considered energy range.

Also, one notes that the ratios of the "indirectly determined" astrophysical S factors, $S_{1\ 3/2}(0)$ and $S_{1\ 1/2}(0)$, for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction populating to the ground and first excited states obtained in the present work to those for the mirror $t(\alpha, \gamma){}^7\text{Li}$ reaction populating to the ground and first excited states deduced in Ref.[22] are equal to $R_S^{(g.s.)}=6.4\pm 0.8$ and $R_S^{(exc)}=6.1\pm 0.7$, respectively. These values are in a good agreement with those of $R_S^{(g.s.)}=6.6$ and $R_S^{(exc)}=5.9$ deduced in Ref.[38] within the microscopic cluster model. This result also confirms directly our estimation for the ratio $R_{C;j_f}$ obtained above since the ANC's for $t + \alpha \rightarrow {}^7\text{Li}(\text{g.s.})$ and $t + \alpha \rightarrow {}^7\text{Li}(0.478 \text{ MeV})$ as well as the ANC's for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ determine the astrophysical S factors for the $t(\alpha, \gamma){}^7\text{Li}$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reactions at zero energies and, consequently, the ratios $R_S^{(g.s.)}$ and $R_S^{(exc)}$ are proportional to $R_{C;3/2}$ and $R_{C;1/2}$, respectively.

Fig. 6d shows a comparison between the branching ratio $R^{exp}(E)$ obtained in the present work (the opened triangle symbols) and that recommended in Refs.[41] (the filled square symbols), in [7, 9] (the filled circle symbols) and [10] (the filled triangle symbols). The weighted mean $\bar{R}^{exp}$ of the $R^{exp}(E)$ recommended by us is equal to $\bar{R}^{exp}=0.41\pm 0.01$. As it is seen from Fig.6d, the branching ratio obtained in the present work and in [7, 9, 10] is in a good agreement with that recommended in Ref.[41] although the underestimation occurs for the $S_{34}^{exp}(E)$ obtained in Ref.[41]. Such a good agreement between two of the experimental data for the $R^{exp}(E)$ can apparently be explained by the fact that there is a reduction factor in [41], being overall for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ astrophysical S factors. The present result for $\bar{R}^{exp}$ is in a good agreement with those of 0.43 ± 0.02 [41] and 0.43 [16, 42] but is noticeably larger than 0.37 [15] and 0.32 ± 0.01 [43].

Thus, it follows from here that the overall normalization of the astrophysical S factors at extremely low energies for the reactions under consideration is mainly determined by the ANC-values for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$. The found ANC's can predict the separated experimental astrophysical S factors for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ reactions at low experimentally acceptable energy regions ($126.5 \leq E \leq 951 \text{ keV}$) obtained by using the total experimental astrophysical S factors [6–9]. Also, they allow us to perform correct extrapolation of the astrophysical S factors for the direct radiative capture ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction at solar energies, including $E=0$.

3 Conclusion

The analysis of the experimental astrophysical S factors, $S_{34}^{exp}(E)$, for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction, which were precisely measured at energies $E=92.9\text{--}1235$ keV [6–10], has been performed within the modified two-body potential approach proposed recently in Ref.[22]. The scrupulous quantitative analysis shows that the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ reaction within the considered energy ranges is mainly peripheral and the parameterization of the direct astrophysical S factors in terms of ANC's for the ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ is adequate to the physics of the peripheral reaction under consideration.

It is demonstrated that the experimental astrophysical S factors of the reaction under consideration measured in the aforesaid energy region can be used as an independent source of getting the information about the ANC's (or NVC's) for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ (or for the virtual decay ${}^7\text{Be} \rightarrow \alpha + {}^3\text{He}$). The weighted means of the ANC's for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$ and NVC's for the virtual decay ${}^7\text{Be} \rightarrow \alpha + {}^3\text{He}$ are obtained. They have to be $(C_{1\ 3/2}^{exp})^2=23.2\pm 1.4$ fm⁻¹ and $(C_{1\ 1/2}^{exp})^2=15.7\pm 1.0$ fm⁻¹ for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(\text{g.s})$ and ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}(0.429\text{ MeV})$, respectively. The corresponding values of the NVC's are $|G_{1\ 3/2}|^2=1.11\pm 0.07$ fm and $|G_{1\ 1/2}|^2=0.750\pm 0.050$ fm. The uncertainty in the ANC (NVC)-values involves the experimental errors for the experimental astrophysical S factors (including that for the activation and the prompt method of measurement) and that of the used approach. Also, the values of ANC's were used for getting the information about the α -particle spectroscopic factors for the mirror (${}^7\text{Li}{}^7\text{Be}$)-pair.

The obtained values of the ANC's were also used for obtaining the experimental ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ astrophysical S factors for capture to the ground and first excited states, the branching ratio at the nine experimental points of energy E (within the interval $E=93.3\text{--}951.0$ keV) [6]–[9] and for an extrapolation of astrophysical S factors at energies less than 90 keV, including $E=0$. In particular, for the weighted mean of the branching ratio $\bar{R}^{exp}$ and the total astrophysical S factor $S_{34}(0)$ the values of $\bar{R}^{exp}=0.41\pm 0.01$ and $S_{34}(0)=0.610\pm 0.037$ keV b have been obtained, respectively. The latter is in an agreement with that $S_{34}(0)=0.56$ keV [13] obtained within the microscopical single-channel ($\alpha + {}^3\text{He}$) cluster model, $S_{34}(0)=0.609$ keV b [12] and $S_{34}(0)=0.621$ keV b [37] obtained within the ($\alpha + {}^3\text{He}$)-channel of version of the resonating-group method, but it is noticeably larger than the result of $S_{34}(0)=0.516$ keV [16] obtained within the standard two-body ($\alpha + {}^3\text{He}$) potential by using $\alpha{}^3\text{He}$ potential deduced by a double-folding procedure.

Acknowledgments

The authors are deeply grateful to L.D. Blokhintsev and A.M. Mukhamedzhanov for discussions and general encouragement. The authors thank also D.Bemmerer for providing the experimental results of the updated data analysis. The work has been supported by The Academy of Science of The Republic of Uzbekistan (Grant No.FA-F2-F077).

References

- [1] J. N. Bahcall, A. M. Serenelli, and S. Basu, *Astrophys. J.* **621**, L85 (2005).
- [2] J. N. Bahcall, W.F. Huebner, S. H. Lubow, P. D. Parker, and R. K. Ulrich, *Rev.Mod.Phys.* **54**, 767 (1982).
- [3] J. N. Bahcall, and M. H. Pinsonneault, *Rev.Mod.Phys.* **64**, 781 (1992).
- [4] E.G. Adelberger, S.M.Austin, J.N. Bahcall, A.B.Balantekin, G.Bogaert, L.S.Brown, L.Buchmann, F.E.Cecil, A.E.Champagne, L. de Braekeleer, Ch.A. Duba, S.R. Elliott, S.J. Freedman, M. Gai, G.Goldring, Ch. R. Gould, A. Gruzinov, W.C. Haxton, K.M. Heeger, and E.Henley, *Rev. Mod. Phys.* **70**, 1265 (1998).
- [5] C.Angulo, M.Arnoult, M.Rayet, P.Descouvemont, D.Baye, C.Leclercq-Willain, A.Coc, S.Barhoumi, P.Aguer, C.Rolfs, R.Kunz, J.W. Hammer, A.Mayer, T. Paradelis, S.Kossionides, C.Chronidou, K.Spyrou, S.Degl’Innocenti, G. Fiorentini, B.Ricci, S.Zavatarelli, C.Providencia, H.Wolters, J.Soaes, C.Grama, J.Rahighi, A.Shotter, M.L.Rachti, *Nucl. Phys. A* **656**, 3 (1999).
- [6] B. S. Nara Singh, M. Hass, Y. Nir-El, and G. Haquin, *Phys.Rev.Lett.* **93**, 262503 (2004).
- [7] D. Bemmerer, F. Confortola, H. Costantini, A. Formicola, Gy. Gyürky, R. Bonetti, C. Brogini, P. Corvisiero, Z. Elekes, Zs. Fülöp, G. Gervino, A. Guglielmetti, C. Gustavino, G. Imbriani, M. Junker, M. Laubenstein, A. Lemut, B. Limata, V.Loza, M. Marta, R. Menegazzo, P. Prati, V. Roca, C. Rolfs, C. Rossi Alvarez, E. Somorjai, O. Straniero, F. Strieder, F. Terrasi, and H.P. Trautvetter (The LUNA Collaboration), *Phys.Rev.Lett.* **97**, 122502 (2006); private communication.
- [8] Gy. Gyürky, F. Confortola, H. Costantini, A. Formicola, D. Bemmerer, R. Bonetti, C. Brogini, P. Corvisiero, Z. Elekes, Zs. Fülöp, G. Gervino, A. Guglielmetti, C. Gustavino, G. Imbriani, M. Junker, M. Laubenstein, A. Lemut, B. Limata, V.Loza, M. Marta, R. Menegazzo, P. Prati, V. Roca, C. Rolfs, C. Rossi Alvarez, E. Somorjai, O. Straniero, F. Strieder, F. Terrasi, and H.P. Trautvetter (The LUNA Collaboration), *Phys.Rev. C* **75**, 035805 (2007) .
- [9] F. Confortola, D. Bemmerer, H. Costantini, A. Formicola, Gy. Gyürky, P. Bezzon, R. Bonetti, C. Brogini, P. Corvisiero, Z. Elekes, Zs. Fülöp, G. Gervino, A. Guglielmetti, C. Gustavino, G. Imbriani, M. Junker, M. Laubenstein, A. Lemut, B. Limata, V.Loza, M. Marta, R. Menegazzo, P. Prati, V. Roca, C. Rolfs, C. Rossi Alvarez, E. Somorjai, O. Straniero, F. Strieder, F. Terrasi, and H.P. Trautvetter (The LUNA Collaboration), *Phys.Rev. C* **75**, 065803 (2007).
- [10] T.A.D. Brown, C. Bordeanu, R.F. Snover, D.W. Storm, D. Melconian, A.L. Sallaska, S.K.L. Sjeue, S. Triambak, *Phys.Rev. C* **76**, 055801 (2007).

- [11] P.Descouvemont, A. Adahchour, C. Angulo, A. Coc, E. Vangioni-Flam. Atomic data and Nuclear data Tables. **88**, 203 (2004).
- [12] T. Kajino, Nucl.Phys. A **460**, 559 (1986).
- [13] K. Langanke, Nucl. Phys. A **457**, 351 (1986).
- [14] A. Csóto and K. Langanke, Few-body Systems. **29**, 121 (2000).
- [15] K.M. Nollett, Phys. Rev. C **63**, 054002 (2001).
- [16] P. Mohr, H. Abele, R. Zwiebel, G. Staudt, H. Krauss, H. Oberhummer, A. Denker, J.W. Hammer and F. Wolf, Phys. Rev. C **48**, 1420 (1993).
- [17] S.B. Dubovitchenko, A.V. Dzhazairov - Kakhramanov, Yad. Fiz. **58**, 635 (1995) [Phys. At. Nucl. **58**, 579 (1995)]. **48**, 1664 (1982); Nucl.Phys. A **419**, 115 (1984).
- [18] R.F. Christy and I. Duck, Nucl.Phys. **24**, 89 (1961).
- [19] S.B. Igamov, T.M. Tursunmuratov, and R. Yarmukhamedov, Yad.Fiz. **60**, 1252 (1997) [Phys. At. Nucl. **60**, 1126 (1997)].
- [20] S.B. Igamov and R. Yarmukhamedov, Yad. Fiz. **58**, 1400 (1995)[Phys. At. Nucl. **58**, 1317 (1995)].
- [21] A.M. Mukhamedzhanov, R.P. Schmitt, R.E. Tribble, and A. Sattarov, Phys. Rev. C **52**, 3483 (1995).
- [22] S.B. Igamov, and R. Yarmukhamedov, Nucl.Phys.A **781**, 247 (2007).
- [23] L.D. Blokhintsev, I.Borbely, E.I. Dolinskii, Fiz.Elem.Chastits At. Yadra. **8**, 1189 (1977)[Sov. J. Part. Nucl. **8**, 485 (1977)].
- [24] L.D. Blokhintsev and V.O. Yeromenko, Yad.Fiz. **71**, 1219 (2008) [Phys. At. Nucl. **71**, 1126 (2008)].
- [25] J.L. Osborne, C. A. Barnes, R.W. Ravanagh, R.M. Kremer, G.J. Mathews, J.L. Zyskind, P.D. Parker, and A.J. Howard, Phys.Rev.Lett.
- [26] R.G.H. Robertson, P. Dyer, R.A. Warner, R.C. Melin, T.J. Bowles, A.B. McDonald, G.C. Ball, W.G. Davies, E.D. Earle, Phys.Rev.Lett. **47**, 1867 (1981).
- [27] K.H. Kim, M.H. Park, B.T. Kim, Phys.Rev. C **35** 363 (1987).
- [28] S.B. Igamov and R. Yarmukhamedov, Phys. At. Nucl. **71**, 1740 (2008)].
- [29] S.V. Artemov, S.B. Igamov, K.I. Tursunmakhatov, and R. Yarmukhamedov, Izv. RAN: Seriya Fizicheskaya, **73**, 176 (2009)[Bull.RAS:Physics,**73**, 165 (2009)].

- [30] S. A. Goncharov, J. Dobesh, E. I. Dolinskii, A. M. Mukhamedzhanov and J. Cejpek, *Yad. Fiz.***35**, 662 (1982)[*Sov. J. Nucl. Phys.* **35**, 383 (1982)].
- [31] V.G. Neudatchin, V.I. Kukulin, A.N. Boyarkina and V.D. Korennoy, *Lett.Nuovo Cim.***5**, 834 (1972).
- [32] V.I. Kukulin, V.G. Neudatchin, and Yu. F. Smirnov, *Nucl.Phys. A* **245**, 429 (1975).
- [33] V.I. Kukulin, V.G. Neudatchin, I.T. Obukhovskiy and Yu.F. Smirnov. "Clusters as Subsystems in Light Nuclei". In *Clustering Phenomena in nuclei*, Vol.3, eds. K. Wildermuth and P. Kramer (Vieweg , Braunschweig, 1983)p.1.
- [34] S.B. Dubovichenko and M.A. Zhusupov, *Izv. Akad. Nauk Kaz.SSR, Ser. Fiz.-Mat.* **4**, 44 (1983); *Yad. Fiz.* **39**, 1378 (1984) [*Sov. J. Nucl. Phys.***39**, 870 (1984).]
- [35] A.L.C. Barnard, C. M. Jones, and G.C. Phillips, *Nucl.Phys.* **50**, 629 (1964).
- [36] C. Rolfs, *Nucl.Phys. A* **217**, 29(1973).
- [37] H. Walliser, H. Kanada, and Y.C. Tang, *Nucl.Phys. A* **419**, 133 (1984).
- [38] N.K. Timofeyuk, P. Descouvemont, R.C. Johnson, *Phys.Rev. C* **75**, 034302 (2007).
- [39] D. Kurath and D. J. Millener, *Nucl.Phys.A* **238**, 269 (1975).
- [40] C.R. Brune, W.H. Geist, R.W. Kavanagh, K.D. Veal, *Phys.Rev.Lett.* **83**, (1999) 4025.
- [41] H.Kräwinkel, H.W. Becker, L. Buchmann, J. Görres, K.U. Kettner, W.E. Kieser, R. Santo, P. Schmalbrock, H.P. Trautvetter, A. Vlieks, C. Rolfs, J.W. Hammer, R.E. Azuma, W.S. Rodney, *Z.Phys. A* **304**, 307 (1982).
- [42] T. Altmeyer, E. Kolbe, T. Warmann, K. Langanke, H.J. Assenbaum, *Z.Phys. A* **330**, 277 (1988).
- [43] U. Schröder, A. Redder, C. Rolfs, R.E. Azuma, L. Buchmann, C. Campbell, J.D. King, T.R. Donoghue, *Phys.Lett. B* **192**, 55 (1987).




Figure 1: The dependence of $\mathcal{R}_{l_f j_f}(E, C_{l_f j_f}^{(sp)})$ as a function of the single-particle ANC, $C_{l_f j_f}^{(sp)}$, for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ $((l_f, j_f) = (1, 3/2))$ and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ $((l_f, j_f) = (1, 1/2))$ reactions at different energies E .




Figure 2: The energy dependence of the $\alpha^3\text{He}$ -elastic scattering phase shifts for different partial waves. The experimental data are from [35]. The bands are our calculated data. The width of the bands for fixed energies corresponds to the variation of the parameters R and a of the adopted Woods–Saxon potential within the intervals of $R=1.62$ to 1.98 fm and $a=0.63$ to 0.77 fm.




Figure 3: The dependence of the ANC's $C_{l_f j_f}$ (upper band) and the spectroscopic factors $Z_{l_f j_f}$ (lower band) on the single-particle ANC $C^{(sp)}_{l_i j_i}$ for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ (the left column, $(l_f, j_f) = (1, 3/2)$) and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(0.429 \text{ MeV})$ (the right column, $(l_f, j_f) = (1, 1/2)$) reactions at different energies E .




Figure 4: The integrand of the radial integral (10) of Ref.[22] for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ reaction at energies $E=0.1056$ and 0.1477 MeV for different sets of $(R; a)$ -pairs: $(1.62 \text{ fm}; 0.63 \text{ fm})$ (dashed line), $(1.80 \text{ fm}; 0.70 \text{ fm})$ (solid line) and $(1.98 \text{ fm}; 0.77 \text{ fm})$ (dotted line).




Figure 5: The values of the ANC's, $C_{1 3/2}^2$ and $C_{1 1/2}^2$, for $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(\text{g.s.})$ (a) and $\alpha + {}^3\text{He} \rightarrow {}^7\text{Be}(0.429 \text{ MeV})$ (b) for each energy E experimental point. The symbols are data obtained from all experimental astrophysical S factors. The opened triangle and cycle symbols (filled star(the activation) and square(the prompt) symbols) are data obtained by using the total (separated) experimental astrophysical S factors from [6] and [7, 9] (from [10]), respectively, while filled circle symbols are data obtained from the separated experimental astrophysical S factors from Refs.[7–9]. The solid lines present our results for the weighted means. Everywhere the width of each of the band is the weighted uncertainty.




Figure 6: The astrophysical S factors for the ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}(\text{g.s.})$ ($(l_f, j_f) = (1, 3/2)$) (a), ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ (0.429 MeV) ($(l_f, j_f) = (1, 1/2)$) (b) and ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ (0.429 MeV) ($(l_f, j_f) = (1 3/2) + (1 1/2)$) (c) reactions as well as the branching ratio (d). In (a) and (b): the opened diamond and triangle symbols are our result separated from the total experimental astrophysical S factors of Refs.[7-9] and [6], respectively; the filled circle symbols (filled star and square symbols) are experimental data of Ref.[9] (Ref.[10], the activation and the prompt method, respectively); the opened circle symbols are our results of the extrapolation; in (c), the symbols are data of all experiments [6]–[10] and the present work (the open circle symbols). The solid lines present our calculations performed with the standard values of geometric parameters $R=1.80$ fm and $a=0.70$ fm both for the bound ($\alpha + {}^3\text{He}$) state and for $\alpha^3\text{He}$ -scattering state. The dashed line is the result of [16]. In (d): the filled circle, triangle and square symbols are experimental data taken from Refs.[7, 9], [10] and [41], respectively, and the opened triangle symbols are our results. The straight line and width of band are our results for the weighted mean and its uncertainty, respectively.

Table 1: The "indirectly determined" values of the asymptotic normalization constants $((C_{13/2}^{exp})^2$ and $(C_{11/2}^{exp})^2$) for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$, the experimental astrophysical S factors ($S_{1j_f}^{exp}$ and $S_{34}^{exp}(E)$) and branching ratio ($R^{exp}(E)$) at different energies E .

E (keV)	$(C_{1j_f}^{exp})^2$ (fm $^{-1}$)		$S_{1j_f}^{exp}$ (keV b)		$S_{34}^{exp}(E)$ (keV b)	$R^{exp}(E)$
	$j_f=3/2$	$j_f=1/2$	$j_f=3/2$	$j_f=1/2$		
92.9 ^{*)}	22.0 $\pm$ 1.8	14.0 $\pm$ 1.2	0.387 $\pm$ 0.031[7, 9]	0.147 $\pm$ 0.012[7, 9]	0.534 $\pm$ 0.023[7, 9]	0.380 $\pm$ 0.030[7, 9]
93.3 ^{**)}	21.4 $\pm$ 1.4	14.7 $\pm$ 0.9	0.374 $\pm$ 0.020	0.153 $\pm$ 0.009	0.527 $\pm$ 0.027[9]	0.409 $\pm$ 0.03
105.6 ^{*)}	21.0 $\pm$ 1.8	14.6 $\pm$ 1.2	0.365 $\pm$ 0.030[7, 8]	0.151 $\pm$ 0.012[7, 8]	0.516 $\pm$ 0.031[7, 8]	0.415 $\pm$ 0.029[7, 9]
106.1 ^{**)}	21.2 $\pm$ 1.3	14.6 $\pm$ 0.9	0.368 $\pm$ 0.020	0.150 $\pm$ 0.009	0.518 $\pm$ 0.024[9]	0.408 $\pm$ 0.020
126.5 ^{*)}	21.2 $\pm$ 0.9	14.6 $\pm$ 0.6	0.366 $\pm$ 0.023	0.148 $\pm$ 0.009	0.514 $\pm$ 0.019[7, 8]	0.404 $\pm$ 0.020
147.7 ^{*)}	20.8 $\pm$ 1.1	14.6 $\pm$ 0.7	0.352 $\pm$ 0.017[7]	0.147 $\pm$ 0.007[7]	0.499 $\pm$ 0.017[7, 8]	0.417 $\pm$ 0.020[7]
168.9 ^{*)}	20.6 $\pm$ 0.8	14.1 $\pm$ 0.6	0.343 $\pm$ 0.010	0.139 $\pm$ 0.006	0.482 $\pm$ 0.017[7, 8]	0.405 $\pm$ 0.020
170.1 ^{**)}	21.9 $\pm$ 1.2	15.0 $\pm$ 0.8	0.362 $\pm$ 0.020	0.148 $\pm$ 0.008	0.510 $\pm$ 0.021[9]	0.409 $\pm$ 0.030
420.0 ^{*)}	21.4 $\pm$ 1.7	14.7 $\pm$ 1.1	0.297 $\pm$ 0.020	0.123 $\pm$ 0.009	0.420 $\pm$ 0.030[6]	0.414 $\pm$ 0.050
506.0 ^{*)}	20.9 $\pm$ 1.9	14.3 $\pm$ 1.3	0.266 $\pm$ 0.020	0.113 $\pm$ 0.010	0.379 $\pm$ 0.030[6]	0.424 $\pm$ 0.050
615.0 ^{*)}	21.5 $\pm$ 1.4	14.7 $\pm$ 0.9	0.254 $\pm$ 0.020	0.108 $\pm$ 0.006	0.362 $\pm$ 0.020[6]	0.425 $\pm$ 0.040
951.0 ^{*)}	22.7 $\pm$ 1.2	15.6 $\pm$ 0.8	0.220 $\pm$ 0.010	0.096 $\pm$ 0.005	0.316 $\pm$ 0.010[6]	0.436 $\pm$ 0.030

^{*)} the activation

^{**)} the prompt method

Table 2: The weighted means of the ANC-values $(C^{\text{exp}})^2$ for ${}^3\text{He} + \alpha \rightarrow {}^7\text{Be}$, NVC's $|G|_{\text{exp}}^2$ and the calculated values of $S_{34}(E)$ at energies $E=0$ and 23 keV . The the second and third lines (the fifth and sixth lines) correspond to the results obtained by means of the analysis of data from works pointed out in the first column, and the list line corresponds to that obtained by using data from all experiments [6–10]. The figures parentetical are the weighted means obtained from corresponding two ones given for the activation and the prompt method.

Experimental data	$(C_{13/2}^{\text{exp}})^2, \text{ fm}^{-1}$	$ G_{13/2} _{\text{exp}}^2, \text{ fm}$	$(C_{11/2}^{\text{exp}})^2, \text{ fm}^{-1}$	$ G_{11/2} _{\text{exp}}^2, \text{ fm}$	$S_{34}(0), \text{ keV b}$	$S_{34}(23 \text{ keV}), \text{ keV b}$
[10](the activation)	24.0 ± 0.8	1.14 ± 0.04	16.3 ± 0.6	0.770 ± 0.030	0.630 ± 0.022	0.619 ± 0.022
[10](the prompt method)	23.7 ± 1.0	1.13 ± 0.05	16.1 ± 0.9	0.760 ± 0.050	0.624 ± 0.029 (0.628 ± 0.018) 0.596 ± 0.021 [10]	0.612 ± 0.019
[6]–[9](the activation)	21.2 ± 0.7	1.01 ± 0.03	14.6 ± 0.4	0.690 ± 0.020	0.560 ± 0.010	0.551 ± 0.009
[9](the prompt method)	21.5 ± 0.3	1.02 ± 0.01	14.8 ± 0.2	0.700 ± 0.010	0.568 ± 0.004 (0.567 ± 0.004) 0.560 ± 0.017 [9]	0.558 ± 0.004
[6]–[10]	23.2 ± 1.4	1.11 ± 0.07	15.7 ± 1.0	0.750 ± 0.050	0.610 ± 0.037	0.599 ± 0.036